

Pancyclicity when each cycle must pass exactly k Hamilton cycle chords

Fatima Affif Chaouche*, Carrie Rutherford†, Robin Whitty†

December 18, 2012

Abstract It is known that $\Theta(\log n)$ chords must be added to an n -cycle to produce a pancyclic graph; for vertex pancyclicity, where every vertex belongs to a cycle of every length, $\Theta(n)$ chords are required. A possibly ‘intermediate’ variation is the following: given k , $1 \leq k \leq n$, how many chords must be added to ensure that there exist cycles of every length each of which passes exactly k chords? For fixed k , we establish a lower bound of $\Omega(n^{1/k})$ on the growth rate.

Keywords: extremal graph theory, pancyclic graph, Hamilton cycle.

A simple graph G on n vertices is *pancyclic* if it has cycles of every length l , $3 \leq l \leq n$. The study of these graphs was initiated by Bondy’s observation [1, 2] that, for non-bipartite graphs, sufficient conditions for hamiltonicity can also be sufficient for pancyclicity. A specific example is the Ore-type hamiltonicity condition, that for 2-connected G , $d(x) + d(y) \geq n$ for all nonadjacent $x, y \in V(G)$. Bondy [1] showed this condition also implies that either G is pancyclic or that n is even and G is $K_{n/2, n/2}$.

That Bondy’s theorem gives a sufficient but not necessary condition is easy to see: label the vertices around the cycle C_6 , in order, as $v_1, \dots, v_6$. Add chords v_1v_3 and v_2v_6 ; the result fails to satisfy the hypothesis of Bondy’s theorem but is nonetheless a pancyclic graph. It is the role of chords in pancyclicity that is explored in this note. In our C_6 with two added chords, the 3 and 5-cycles must both use one chord; the 4-cycle must use two, while there are Hamilton cycles using zero or two chords. If a third chord v_1v_4 is added then we can find cycles of all lengths each using exactly two chords.

In general, we may distinguish, in a pancyclic graph G , a Hamilton cycle C ; then the remaining edges of G form chords of C . We can then ask, given $k \leq l \leq n$ if, relative to C , a cycle of length l exists which uses exactly k chords. This suggests a k -chord analog of pancyclicity: do all possible cycle lengths occur when cycles must use exactly k -chords of a suitably chosen Hamilton cycle?

Obviously some restrictions apply: no Hamilton cycle can use exactly one chord of another Hamilton cycle; we cannot form an $(n - 1)$ -chord Hamilton cycle in C_n unless $n \geq 6$, etc, etc. To avoid cataloguing special cases, a definition of ‘possessing all k -chord cycles physically possible’ is proposed as follows:

Take a copy of C_n and add a chord between all pairs of non-adjacent vertices to obtain an isomorphic copy of K_n . Denote by $S(n, k)$ the set of lengths of cycles in this graph which pass exactly k chords. Call a graph G on n vertices *k -chord pancyclic* if G contains a Hamilton cycle relative to which, for each integer $l \in S(n, k)$, G contains a cycle of length l passing exactly k chords.

*University of Sciences and Technology Houari Boumediene, Algiers

†London South Bank University

- Example 1** 1. The copy of C_6 , with added chords v_1v_3 , v_2v_6 and v_1v_4 , as described above, was observed to be 2-chord pancyclic. If two further chords, v_2v_4 and v_4v_6 , are added then the result is a 3-chord pancyclic graph, as may be verified by inspection.
2. It is easy to see that $G = K_{n/2, n/2}$ contains two edge-disjoint Hamilton cycles for n even, $n \geq 8$. Since G contains no odd-length cycles, G is k -chord pancyclic precisely for $k = n$. General conditions for disjoint Hamilton cycles in bipartite graphs, e.g. Ferrara et al [5], may be used to extend this example.
3. The graph in (2) is an extreme example of where k -chord pancyclicity does not imply pancyclicity. Clearly for $k \leq 3$ the implication does hold. For $k = 4$ it need not, as demonstrated by the graph in figure 1(a) which is 4-chord pancyclic and has girth 4.
4. The graphs in figure 1(b) and (c) are isomorphic, as indicated by the vertex labelling. The graph is seen to be 1-chord pancyclic using the chords defined in (b) but not using the chords defined in (c). Thus for a given Hamiltonian graph G the property of k -chord pancyclicity may hold relative to one Hamilton cycle but not to another.




Figure 1: (a) a 4-chord pancyclic graph of girth 4; (b) a graph which is 1-chord pancyclic with respect to the given C_9 ; (c) no 4-cycle uses exactly 1 chord of the given C_9 .

A natural question to ask would seem to be: when is pancyclicity enough to guarantee k -chord pancyclicity and vice versa? However example 1(4) suggests that such questions may be difficult to study because the answers may depend on making a suitable choice of Hamilton cycle. In the present paper we are less ambitious: starting with an n -cycle, we construct a k -chord pancyclic graph by adding chords to this cycle. We offer a bound on the minimum number of chords necessary to guarantee k -chord pancyclicity in this sense. The same approach may yield simple bounds for ordinary pancyclicity; the following is stated without proof in [1]:

Theorem 2 In a pancyclic graph G on n vertices the number of edges is not less than $n - 1 + \log_2(n - 1)$. $\square$

It is worthwhile to supply a proof of theorem 2 since it follows immediately from the upper bound in the following lemma, which we refer to later on:

Lemma 3 Suppose p chords are added to C_n , $n \geq 3$. Then the number $N(n, p)$ of cycles in the resulting graph satisfies

$$\binom{p+2}{2} \leq N(n, p) \leq 2^{p+1} - 1.$$

Proof. Embed C_n convexly in the plane. Suppose the chords added to C_n are, in order of inclusion, $e_1, e_2, \dots, e_p$. Say that e_i intersects e_j if these edges cross each other when added to the embedding of C_n . Let n_i be the number of new cycles obtained with e_i is added. Then n_i satisfies:

1. $n_i \geq i + 1$, the minimum occurring if and only if the e_j are pairwise non-intersecting for, $j \leq i$;
2. $n_i \leq 2^i$, the maximum occurring if and only if e_i intersects with e_j for all $j < i$, giving $n_i = \sum_{j=0}^i \binom{i}{j}$.

Now $1 + \sum_{i=1}^p (i + 1) \leq 1 + \sum_{i=1}^p n_i \leq 1 + \sum_{i=1}^p 2^i$ and the result follows. $\square$

The bound of theorem 2 works well for small values of n . The minimum number of edges in a pancyclic graph for $3 \leq n \leq 22$ is given in [6]. For $3 \leq n \leq 14$ the lower bound in theorem 2 is achieved; however, it can be seen that, for $n = 15, 16$, we must add four chords to C_n to achieve pancyclicity while the argument in the proof of lemma 3 can only account for three.

As regards an upper bound on the number of chords required for pancyclicity, [1] also gives an $O(\log n)$ bound, again without a construction. A $\log n$ construction is given by Sridharan [7]. We may summarise by saying that pancyclicity is achieved by adding $\Theta(\log n)$ chords to C_n .

In contrast, *vertex pancyclicity*, in which each vertex must lie on a cycle of every length, has been shown by Broersma [3] to require $\Theta(n)$ edges to be added to C_n . Our question is: where between $\log n$ and n does k -chord pancyclicity lie?

Let us for the moment restrict to $k \geq 3$. Suppose we add p chords to C_n , $3 \leq k \leq p \leq \binom{n}{2} - n$, necessarily requiring $n \geq 6$. Suppose that these p added chords include a k -cycle (obviously required for k -chord pancyclicity). We will use $K(k, p)$, defined for $k \geq 3$, to denote the maximum number of k -chord cycles that can be created in the resulting graph. Then $1 \leq K(k, p)$ by definition and $K(k, p) \leq 2^{p+1} - 1$ by lemma 3. By lowering this upper bound we can increase the lower bound on the number p of chords required for k -chord pancyclicity.

Theorem 4 $K(k, p) \leq \binom{p}{k} + k \binom{p-k}{k-1} + \binom{p-k}{k}$.

We will use the following Lemma to prove theorem 4:

Lemma 5 Suppose that a set X of chords is added to C_n . In the resulting graph the maximum number of cycles passing all edges in X is

$$\begin{cases} 1 & \text{if } X \text{ contains adjacent chords} \\ 2 & \text{if no two chords of } X \text{ are adjacent} \end{cases}$$

Proof. Let G be the graph resulting from adding the chords of X to C_n . We may assume without loss of generality that G has no vertices of degree 2, since such vertices may be contracted out. For a given cycle in G passing all chords of X , let H denote the intersection of this cycle with the C_n . Then H consists of isolated vertices and disjoint edges, and H is completely determined once any of these vertices or edges is fixed. If two chords are adjacent this fixes an isolated vertex of H ; if no two chords are adjacent then there is a maximum of two ways in which a single edge of H may be fixed. $\square$

Proof of theorem 4. By definition of $K(k, p)$ we must use a set, say S , of k chords to create a k -cycle. We add new chords to S , one by one. On adding the r -th additional chord, $1 \leq r \leq p-k$, we ask how many k -chord cycles use this chord. For any such a cycle the previous $r-1$ chords will be split between S and non- S chords: with i chords from S being used, $0 \leq i \leq k-1$, this can happen in

$$\binom{k}{i} \binom{r-1}{k-i-1}$$

ways. Since $i > 1$ forces two adjacent chords in S to be used, summing over i , according to lemma 5, and then over r gives

$$K(k, p) \leq 1 + \sum_{r=1}^{p-k} \left(2 \sum_{i=0}^1 \binom{k}{i} \binom{r-1}{k-i-1} + \sum_{i=2}^{k-1} \binom{k}{i} \binom{r-1}{k-i-1} \right).$$

This simplifies (e.g. using symbolic algebra software such as Maple) to give the result. $\square$

Corollary 6 *Let $k \leq n$ be positive integers, with $n \geq 6$ and $k \geq 3$. Let C be a copy of C_n . The number p of chords which must be added to C in order that the resulting graph is k -chord pancyclic, relative to C , is not less than the largest root of the following polynomial in p :*

$$\Pi(p; n, k) = \binom{p}{k} + k \binom{p-k}{k-1} + \binom{p-k}{k} - n + k - 1.$$

$\square$

We finally extend our analysis to include the cases $k = 1, 2$:

Corollary 7 *Let $n \geq 4$ be a positive integer, and let C be a copy of C_n . The number p of chords which must be added to C in order that the resulting graph is k -chord pancyclic, relative to C , for $k \geq 1$ fixed, is of order $\Omega(n^{1/k})$.*

Proof. It is easy to see that, for $k = 1$, exactly $p = \left\lceil \frac{n-3}{2} \right\rceil$ chords are required to make C_n 1-chord pancyclic.

For $k = 2$ an analysis similar to that used in the proof of theorem 4 shows that the number of 2-chord cycles which may be created by adding p chords to C_n is at most $p^2 - p - 1$. So to have 2-chord cycles of all lengths from 3 to n we require $p^2 - p - 1 \geq n - 2$. In this case we can solve explicitly to get the bound $p \geq \frac{1}{2} (1 + \sqrt{4n-3})$.

Now suppose $k \geq 3$. In order to have all k -chord cycles of all lengths between k and n we must have

$$n - k + 1 \leq \binom{p}{k} + k \binom{p-k}{k-1} + \binom{p-k}{k} \leq f(k)p^k,$$

for some function $f(k)$. Therefore $p^k \geq (n - k + 1)/f(k)$ so, for k fixed, $p = \Omega(n^{1/k})$. $\square$

We are suggesting that k -chord pancyclicity may be ‘intermediate’ between pancyclicity and vertex pancyclicity in the sense that the number of chords it requires to be added to C_n may lie between $\log n$ and n . Thus far we have only a lower bound in support of our suggestion. Moreover, a comparison of the growth orders, $\Omega(\log n)$ as opposed to $\Omega(n^{1/k})$, suggests that this is very much a ‘for large n ’ type result. The equation $\ln n = n^{1/k}$ has two positive real solutions for $k \geq 3$, given in terms of the two real branches of the Lambert W function [4]. In particular $\ln n$ exceeds $n^{1/k}$ for $n > e^{-kW_{-1}(-1/k)}$, and this bound grows very fast with k : at

least two orders of magnitude per unit increase! To give a specific example, for 10-chord pancyclicity, the log bound exceeds the 10-th root bound until the number of vertices exceeds about 3.4×10^{15} . Until then, so far as our analysis goes, we might expect ‘most’ pancyclic graphs to be 10-chord pancyclic. However we suggest that, in the long term, a guarantee of this implication, analogous to hamiltonicity guaranteeing pancyclicity, will not be found.

References

- [1] Bondy, J.A., “Pancyclic graphs I”, *J. Combinatorial Theory B*, 11 (1), 1971, 80-84.
- [2] Bondy, J.A., “Pancyclic graphs: recent results, infinite and finite sets”, in : *Colloq. Math. Soc. János Bolyai*, Keszthely, Hungary, 1973, 181-187.
- [3] Broersma, H.J., “A note on the minimum size of a vertex pancyclic graph”, *Discrete Math.*, 164, 1997, 29–32.
- [4] Corless, R., Gonnet, G., Hare, D., Jeffrey, D. and Knuth, D., “On the Lambert W function”, *Advances in Computational Mathematics*, 5, 1996, 329-359.
- [5] Ferrara, M., Gould, R., Tansey, G. and Whalen, T., ”Disjoint hamiltonian cycles in bipartite graphs”, *Discrete Mathematics*, 309 (12) 2009, 3811–3820.
- [6] George, J.C., Marr, A. and Wallis, W., “Minimal pancyclic graphs”, *J. Combinat. Math. and Combinat. Comput.*, to appear.
- [7] Sridharan, M. R., On an extremal problem concerning pancyclic graphs, *J. Math. Phys. Sci.*, 12, 1978, 297-306.